

GCE Examinations
Advanced Subsidiary / Advanced Level

Statistics
Module S1

Paper J

MARKING GUIDE

This guide is intended to be as helpful as possible to teachers by providing concise solutions and indicating how marks should be awarded. There are obviously alternative methods that would also gain full marks.

Method marks (M) are awarded for knowing and using a method.

Accuracy marks (A) can only be awarded when a correct method has been used.

(B) marks are independent of method marks.



Written by Shaun Armstrong & Chris Huffer


© *Solomon Press*

These sheets may be copied for use solely by the purchaser's institute.

S1 Paper J – Marking Guide

1. (a) cum. freqs: 3, 9, 24, 44, 53, 55
- (i) median = $28^{\text{th}} = 23.5 + \left(\frac{4}{20} \times 2\right) = 23.9$ g M1 A1
- (ii) 33^{rd} percentile = $\frac{33}{100} \times (55 + 1)^{\text{th}} = 18.48^{\text{th}}$ value M1
 $= 21.5 + \left(\frac{9.48}{15} \times 2\right) = 22.8$ g M1 A1
- (b) 24 - 25: class width 2 \rightarrow 1 cm \therefore class width 1 \rightarrow 0.5 cm M1
 freq. den. = $\frac{20}{2} = 10 \rightarrow 20$ cm \therefore freq. den. 1 \rightarrow 2 cm M1
- (i) 20 - 21: class width 2 \therefore width 1 cm A1
 freq. den. = $\frac{6}{2} = 3 \therefore$ height 6 cm A1
- (ii) 26 - 29: class width 4 \therefore width 2 cm A1
 freq. den. = $\frac{9}{4} = 2.25 \therefore$ height 4.5 cm A1 **(11)**
-
2. (a) $\sum P(x) = k + \frac{1}{2}k + \frac{1}{3}k + \frac{1}{4}k = \frac{25}{12}k = 1 \therefore k = \frac{12}{25}$ M2 A1
- (b) $\frac{12}{25} + \frac{6}{25} = \frac{18}{25}$ M1 A1
- (c) $\sum xP(x) = \frac{12}{25} + \frac{12}{25} + \frac{12}{25} + \frac{12}{25} = \frac{48}{25}$ M1 A1
- (d) $E(X^2) = \sum x^2P(x) = \frac{12}{25} + \frac{24}{25} + \frac{36}{25} + \frac{48}{25} = \frac{24}{5}$ M1 A1
 $E(X^2 + 2) = \frac{24}{5} + 2 = \frac{34}{5}$ M1 A1 **(11)**
-
3. (a) $P(Z > \frac{165-156}{\sqrt{73}}) = P(Z > 1.05) = 0.1469$ M2 A1
- (b) $1 - (0.5 + 0.1469) = 0.3531$ M1 A1
- (c) $P(14\text{yo} > 165) = P(Z > \frac{165-160}{\sqrt{79}}) = P(Z > 0.56) = 0.2877$ M2 A1
 $P(\text{both} > 165) = 0.1469 \times 0.2877 = 0.0423$ (3sf) M1 A1
- (d) more as e.g. answer to (c) satisfies condition but can also have one less than 165 if the other is sufficiently over 165 B2 **(12)**
-
4. (a) mean = $\frac{427}{20} = 21.35$ minutes M1 A1
 variance = $\frac{11077}{20} - 21.35^2 = 98.0$ minutes² (3sf) M2 A1
- (b) for 2nd sample: $\frac{\sum t}{30} = 18.5 \therefore \sum t = 30 \times 18.5 = 555$ M1
 $\frac{\sum t^2}{30} - 18.5^2 = 8.2^2 \therefore \sum t^2 = 30(8.2^2 + 18.5^2) = 12284.7$ M2 A1
 for combined sample: mean = $\frac{427+555}{50} = 19.6$ minutes (3sf) M1 A1
 variance = $\frac{11077+12284.7}{50} - 19.6^2 = 81.5$ minutes² (3sf) M1 A1 **(13)**
-

5. (a)



B3

(b) (i) $\frac{21}{75} = \frac{7}{25}$

M1 A1

(ii) $\frac{13}{39} = \frac{1}{3}$

M1 A1

(c) (i) $\frac{47}{75} \times \frac{46}{74} \times \frac{45}{73} = 0.240$ (3sf)


M2 A1

(ii) $1 - P(\text{all male}) = 1 - \left(\frac{39}{75} \times \frac{38}{74} \times \frac{37}{73}\right) = 0.865$ (3sf)

M3 A1

(14)

6. (a)



B4

(b) e.g. the first six values lie roughly along a straight line but this changes for the two values above 500 g

B2

(c) $S_{ml} = 39540 - \frac{1550 \times 119}{6} = 8798.33$

M1

$S_{mm} = 552500 - \frac{1550^2}{6} = 152083$

M1

$b = \frac{8798.33}{152083} = 0.05785$

M1 A1

$a = \frac{119}{6} - \left(0.05785 \times \frac{1550}{6}\right) = 4.888$

M1 A1

$\therefore a = 4.89, b = 0.0579$

(d) a is the length of the spring with no mass suspended from it

B1

b is the extra extension for each additional gram suspended from spring

B1

(14)

Total

(75)

